



KEY POINTS

- There is no proof that it is safe to swallow artificial fluoride. (1)
- Boris Johnson, along with 92 other MPs, signed an Early Day Motion to prevent water fluoridation in 2003. (2)
- Over 98% of Europe does not deliberately fluoridate drinking water. (3)
- Fluoride in drinking water and food has been shown to reduce a child's intelligence and increase the likelihood of ADHD. (4)(5)
- Tooth decay does not increase when fluoridation ceases. (6)
- Artificial fluoride is the industrial hazardous waste – hexafluorosilicic acid (HFSA). Its harmful ingredients are listed in British Standard BSEN12175:2013. (7)

SOURCES

- (1) <https://www.fluoridefreepeel.ca/wp-content/uploads/2014/06/Health-Canada-FOI-Response-Letter-June2014.pdf>
- (2) <https://edm.parliament.uk/early-day-motion/22866/fluoridation-of-public-water-supplies>
- (3) https://en.wikipedia.org/wiki/Water_fluoridation_by_country
- (4) <https://fluoridealert.org/news/experts-confirm-extremely-low-levels-of-fluoride-reduce-iq-according-to-the-fluoride-action-network/>
- (5) <https://ehjournal.biomedcentral.com/articles/10.1186/s12940-015-0003-1>
- (6) <https://fluoridealert.org/studies/caries05/>
- (7) British Standards Institute. BSEN 12175:2013
- (8) <https://pubmed.ncbi.nlm.nih.gov/6685423>
- (9) Media Polls. www.ukfffa.org.uk, Resources, Fluoridation Polls and Consultations
- (10) Committee on Toxicity (2002). Consideration of Fluoride in the 1997 Total Diet Survey. <https://cot.food.gov.uk/sites/default/files/cot/fluoride.pdf>
- (11) www.ukfffa.org.uk. Resources, An Infant's Exposure to Fluoride from 5 sources; Overexposure per kg weight.
- (12) www.ukfffa.org.uk. Resources. Graph. Infant Exposure to Fluoride from Spak, C.J., et al (1983). "Fluoride in human milk". Acta Paediatr Scand. 1983 Sep; 72(5):699-701.



**LESS THAN 6 MILLION PEOPLE
IN THE UK DRINK FLUORIDATED
WATER. THE LAST
FLUORIDATION SCHEME
STARTED IN 1987. THAT'S
HOW POPULAR FLUORIDATION
IS(N'T).(9)**

**There is no proof that it is safe to
swallow fluoride.**



EFFORTS HAVE BEEN MADE

No evidence that it's safe

Freedom of Information replies by Health Canada in 2014 and by Public Health England in 2018. Requests were made for high quality proof that swallowed fluoride is safe to drink.

Health Canada confirmed it has no toxicology studies on HFSA, no double-blind randomized clinical trials of fluoridation, etc.

Access to Information Request A-2014-00168 <https://www.fluoridefreepeel.ca/wp-content/uploads/2014/06/Health-Canada-FOI-ResponseLetter-June2014.pdf>

Public Health England could not find any research to prove that swallowing fluoride is safe.

Ref: FoI 27/02/lh/914 dated 20th March 2018.



2021 petition, USA

This year, the case against water fluoridation revolves around a Federal Court Case in the USA where the petitioners are asking the Court to oblige the Environmental Protection Agency (EPA) to process the Petition submitted by several Non-Governmental Agencies in 2016 and strengthened in 2020. The Petition asked the EPA to apply the Toxic Substances Control Act to fluoride. Fluoride is currently a "presumed" developmental neurotoxin this being a gnat's whisker away from "known".



FLUORIDE REDUCES COGNITIVE FUNCTION IN UNBORN CHILDREN

It starts in the womb

There is more than ample evidence that fluoride in the womb and in fluoridated baby formula has the effect of reducing a child's intelligence and it also causes ADHD and sleeplessness amongst teenagers.

In 2002, the UK Government's Committee on Toxicity (CoT) advised that no more than 0.05 mg fluoride per kilo body weight per day should be given to 6-month-old babies. It is known that fluoride accumulates in bones and in teeth so the infant should have been considered just as vulnerable as 6-month-old babies. If we apply the maximum of 0.05 mg fluoride/kilo/bodyweight/day to all babies aged 0 – 12 months, it is obvious that if formula-fed using fluoridated water, they are all over-exposed to fluoride.(10) (11)

The graph below (12) demonstrates that this maximum limit is clearly not acknowledged.



UK fluoride hot spots

Infants below the age of 6 months are overdosing on fluoride when fed with baby formula made up with fluoridated tap water.

In areas where there is natural fluoride such as in Essex where the concentration of natural fluoride varies from 0.4 - 1 mg fluoride/litre and in Hartlepool at 1.3mg fluoride/litre, infants are equally or more challenged.

It's not just plain tap water...

We also have to consider the insult to the developing brains of the unborn child when the mother is fluoridated and/or drinks tea which contains highish levels of fluoride. The topic of reduced intelligence caused by fluoride is dealt with on www.ukfffa.org.uk:

This leaflet is printed and distributed by Fluoride Free Alliance UK. Contact www.ukfffa.org.uk, Home > drop-down menu > Contact us